


Table 15 (continued): 1988-1991 woody vegetation change by land cover by native pasture community (See compilation notes on page 10)

Native Pasture Communities		Rate of woody vegetation change (,000ha per year)								% Wooded vegetation cover 1988	% Total clearing in QLD
Name	Area (,000ha)	New woody regrowth	Clearing by Replacement Landcover						Total Cleared		
			Pasture	Crops	Forest	Mining	Infra-structure	Settle-ment			
Bothriochloa-Danthonia (traprock)	328	0.077	3.500	0.000	0.008	0.000	0.070	0.000	3.579	42.89	0.49
Brigalow (northern)	3918	3.666	92.695	2.392	0.029	0.308	0.279	0.038	95.740	45.94	13.12
Brigalow (Roma region)	1622	0.116	7.376	0.446	0.038	0.061	0.029	0.000	7.949	23.04	1.09
Brigalow and Belah (southern)	3226	0.402	21.904	18.187	0.205	0.000	0.112	0.000	40.409	28.04	5.54
Channel Pastures	5534	0.002	0.401	0.000	0.000	0.000	0.003	0.000	0.404	5.83	0.06
Georgina gidgee	1637	0.000	0.021	0.000	0.000	0.000	0.000	0.000	0.022	15.03	0.00
Gidgee (central)	1318	0.240	32.402	0.460	0.000	0.001	0.213	0.000	33.075	51.96	4.53
Gidgee (far west)	611	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	9.19	0.00
Gidgee (southern)	772	0.010	6.041	0.000	0.000	0.000	0.047	0.000	6.088	47.41	0.83
Heath	942	0.000	0.003	0.000	0.000	0.000	0.037	0.000	0.041	91.53	0.01
Littoral and sporobolus	1488	0.069	0.134	0.123	0.000	0.042	0.093	0.144	0.535	37.80	0.07
Mitchell Grass (Ashy Downs)	2556	0.000	0.264	0.000	0.000	0.000	0.038	0.000	0.302	6.26	0.04
Mitchell Grass (Northern Downs)	21963	0.008	36.529	0.000	0.000	0.000	0.223	0.000	36.751	10.37	5.04
Mitchell Grass (Southern Downs)	2553	0.062	9.400	0.062	0.000	0.000	0.061	0.000	9.523	37.43	1.30
Mitchell Grass (Southern Flooded)	875	0.000	0.028	4.182	0.000	0.000	0.209	0.000	4.419	35.56	0.61
Mitchell Grass (Stony Downs)	2143	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	3.48	0.00
Mulga on residuals	8023	0.007	13.059	0.000	0.000	0.001	0.153	0.000	13.213	28.30	1.81
Mulga-Whitewood	637	0.000	0.000	0.000	0.000	0.000	0.020	0.000	0.020	19.52	0.00
Native Sorghum	1013	0.000	0.051	0.000	0.000	0.000	0.079	0.000	0.130	98.23	0.02
Queensland Blue Grass (central)	1574	0.020	16.409	0.926	0.000	0.057	0.126	0.000	17.519	30.39	2.40
Queensland Blue Grass (southern)	940	0.001	0.127	0.029	0.000	0.000	0.000	0.001	0.158	8.62	0.02
Rainforest	1849	0.105	0.665	0.233	0.219	0.005	0.082	0.104	1.307	84.59	0.18
Ribbongrass	847	0.000	0.005	0.000	0.000	0.000	0.079	0.000	0.084	88.70	0.01

Table 15 (continued): 1988-1991 woody vegetation change by land cover by native pasture community *(See compilation notes on page 10)*

Native Pasture Communities		New woody regrowth	Rate of woody vegetation change (,000ha per year) Clearing by Replacement Landcover							% Wooded vegetation cover 1988	% Total clearing in QLD
Name	Area (,000ha)		Pasture	Crops	Forest	Mining	Infra-structure	Settle-ment	Total Cleared		
Schizachrium (flooded plains)	1782	0.022	0.047	0.000	0.000	0.000	0.033	0.000	0.080	57.21	0.01
Schizachrium (hills and plains)	7575	0.000	0.704	0.000	0.000	0.560	1.007	0.001	2.272	95.34	0.31
Soft and hard mulga on red plain	10119	0.133	42.414	0.000	0.000	0.000	1.977	0.000	44.392	62.32	6.08
Spinifex (CQ desert)	4743	0.298	24.372	0.000	0.000	0.000	0.541	0.000	24.913	82.97	3.41
Spinifex (hard on sand dunes)	5458	0.000	0.391	0.000	0.000	0.000	0.044	0.000	0.435	3.71	0.06
Spinifex (hard with eucalypts & acacia)	1987	0.000	3.626	0.000	0.000	0.000	0.058	0.000	3.684	27.99	0.50
Spinifex (Mt Isa highlands)	8895	0.005	0.279	0.000	0.000	0.004	0.430	0.000	0.713	48.83	0.10


Woody Vegetation Change by Regional Vegetation Management Group



© The State of Queensland (Department of Natural Resources and Mines) 2003.

Figure 14: Average annual percentage of woody vegetation cover cleared by regional vegetation management group (1988-1991)

QUEENSLAND
PERCENTAGE OF AREA WOODED
BY REGIONAL VEGETATION
MANAGEMENT GROUP
1988



REGIONAL VEGETATION MANAGEMENT GROUPS

© The State of Queensland (Department of Natural Resources and Mines) 2003.

Figure 15: Woody vegetation cover by regional vegetation management group (1988)

Table 11: 1988-1991 woody vegetation change by land cover by regional vegetation management group (See compilation notes on page 10)

Guideline group			Rate of woody vegetation change (,000ha per year)									
Name	Map Ref. No.	Area (,000ha)	New woody regrowth	Clearing by Replacement Landcover							% Wooded vegetation cover 1988	% Total clearing in QLD
				Grazing	Crop	Forestry	Mining	Infra-structure	Settle-ment	Total Cleared		
Brigalow Belt North	1	1962	0.018	6.161	1.183	0.010	0.135	0.312	0.091	7.891	65.97	1.08
Brigalow Belt South	2	15301	4.625	114.550	43.590	1.061	0.003	2.103	0.002	161.310	50.85	22.09
Cape York	3	12116	0.000	1.049	0.000	0.000	0.564	1.306	0.002	2.921	93.02	0.40
Capricorn-Dawson	4	5313	0.515	28.355	0.189	0.565	0.419	0.604	0.049	30.181	48.20	4.13
Central Qld Coast	5	1458	0.008	2.544	0.695	0.156	0.000	0.976	0.032	4.404	78.95	0.60
CHC	6	23187	0.000	1.915	0.000	0.000	0.001	0.092	0.000	2.008	8.34	0.27
Coast Wide Bay	7	2810	0.655	3.053	1.931	2.117	0.016	0.530	0.439	8.084	62.14	1.11
Desert Uplands North	8	3467	0.219	24.246	0.000	0.000	0.053	0.753	0.000	25.052	84.69	3.43
Desert Uplands South	9	3417	0.292	20.900	0.000	0.000	0.000	0.336	0.000	21.236	76.68	2.91
Einasleigh Uplands North	10	7177	0.000	2.060	0.349	0.000	0.104	0.878	0.016	3.407	88.99	0.47
Einasleigh Uplands South	11	4149	0.000	3.016	0.014	0.000	0.125	0.636	0.000	3.791	85.70	0.52
Gulf Plains North	12	13520	0.037	1.181	0.000	0.000	0.000	1.221	0.000	2.402	77.30	0.33
Gulf Plains South	13	9131	0.000	4.057	0.000	0.000	0.000	0.570	0.000	4.628	34.83	0.63
Inland Burnett	14	2993	0.174	5.844	0.429	0.129	0.022	0.124	0.002	6.550	50.36	0.90
Mulga Land	15	18746	0.440	122.190	0.555	0.011	0.000	2.657	0.001	125.414	56.88	17.17
Nebo - Broadsound	16	2602	4.378	31.992	0.235	0.003	0.187	0.348	0.000	32.766	60.95	4.49
New England Tablelands	17	775	0.101	5.256	0.006	0.049	0.000	0.161	0.001	5.474	46.30	0.75
North West Highlands	18	7202	0.005	0.008	0.000	0.000	0.004	0.121	0.000	0.133	50.73	0.02
Northern Highlands	19	4076	0.752	103.356	0.879	0.026	0.230	0.784	0.000	105.276	68.33	14.42
North-western Mitchell Grass Downs	20	16396	0.000	2.042	0.000	0.000	0.000	0.191	0.000	2.233	7.73	0.31
South East Queensland	21	2490	0.633	2.957	0.323	0.815	0.189	0.280	2.295	6.859	55.52	0.94
South-eastern Mitchell Grass Downs	22	7922	0.042	51.888	0.000	0.000	0.000	0.173	0.000	52.061	23.02	7.13
Southern Highlands	23	5319	0.482	107.956	3.294	0.118	0.371	0.806	0.040	112.586	61.68	15.42
Wet Tropics	24	1993	0.264	0.668	2.130	0.513	0.003	0.200	0.080	3.594	81.31	0.49

SLATS Products

For information on all SLATS products contact:

Program Officer, Geographic Data Services, Department of Natural Resources and Mines.
Phone: (07) 38963187
Fax: (07) 34062762

Or visit the SLATS website: www.nrm.qld.gov.au/slats

Acknowledgements

This work is the combined effort of the SLATS project team which includes: John Armston, Gerry Bisshop, Lindsay Brebber, John Carter, Sel Counter, Tim Danaher, Alisa Eustace, Bruce Goulevitch, David Harris, Beverly Henry, Linda Lawrence, Udaya Senarath, Andrew Stewart and Jim Walls. SLATS is funded by NR&M and a Queensland Treasury Service Enhancement Program.

References

- Accad, A., Neldner, V.J., Wilson, B.A. and Niehus, R.E. (2001) Remnant Vegetation in Queensland: Analysis of Pre-clearing, Remnant 1997-1999 Regional Ecosystem Information. Queensland Herbarium, Environmental Protection Agency, Brisbane.
- Armston, J.D., Danaher, T.J., Goulevitch, B.M. and Byrne, M.I. (2002) Geometric correction of Landsat MSS, TM and ETM+ imagery for mapping woody vegetation cover and change detection in Queensland. In: *Proceedings of the 11th Australasian Remote Sensing and Photogrammetry Conference*, Brisbane, September 2 - 6, 2002.
- Barson, M.M. (1996) Remote Sensing of Agricultural Land Cover for the Australian Continent, In: *Proceedings of the 8th Australasian Remote Sensing Conference*, Canberra 25-29 March 1996, Vol. 2, pp. 122-127.
- Barson, M. M., Randall, L. A. and Bordas, V. (2000) Land Cover Change in Australia. Results of the Collaborative Bureau of Rural Sciences - State agencies' project on remote sensing of land cover change. Bureau of Rural Sciences, Canberra.
- Boulter, S.L., Wilson, J., Westrup, J., Anderson, E.R., Turner, E.J. and Scanlan, J.C. (2000) Native Vegetation Management in Queensland, Background, Science and Values, Department of Natural Resources, Brisbane.
- Burrows, W.H., Henry, B.K., Back, P.V., Hoffmann, M.B., Tait, L.J., Anderson, E.R., Menke, N., Danaher, T., Carter, J.O., and McKeon, G.M. (2002). Growth and carbon stock change in eucalypt woodlands in northeast Australia: ecological and greenhouse sink implications. *Global Change Biology*, 8: 769-784.

Burrows, W.H., Hoffmann, M.B., Compton, J.F., Back, P.V. and Tait, L.J. (2000) Allometric relationships and community biomass estimates for some dominant eucalypts in Central Queensland woodlands, *Australian Journal of Botany*, Vol. 48, No. 6.

Carnahan, J.A. (1988) *Map of Australia Present Vegetation*, Australian Surveying and Land Information Group, Canberra.

Collett, L., Goulevitch, B.M. and Danaher, T.J. (1998) SLATS Radiometric Correction : A Semi Automated Multi Stage Process for the Standardisation of Temporal and Spatial Radiometric Differences. In: *Proceedings of the 9th Australasian Remote Sensing and Photogrammetry Conference*, Sydney, Australia, July 1998.

Danaher, T.J., Xiolaing, W. and Campbell, N.A. (2001). Bi-directional reflectance distribution function approaches to radiometric calibration of Landsat ETM+ imagery. *Proceedings of the IEEE 2001 International Geoscience and Remote Sensing Symposium*, University of NSW, Sydney, Australia, July 2001.

Danaher, T. (2002) An empirical BRDF correction for Landsat TM and ETM+ imagery. In: *Proceedings of the 11th Australasian Remote Sensing and Photogrammetry Conference*, Brisbane, September 2 - 6, 2002.

Danaher, T., Carter, J.O., Brook, K.D. and Dudgeon, G. (1992) Broadscale Vegetation Mapping Using NOAA AVHRR Imagery. In: *Proceedings of the Sixth Australasian Remote Sensing Conference, 2-6 November 1992, Wellington, New Zealand*, Vol. 3, pp. 128-137.

Department of Natural Resources (1997) Interim Report - The Statewide Landcover and Trees Study (SLATS), October 1997. pp. 37.

Department of Natural Resources (1999A) Land Cover Change in South East Queensland 1988-1997, A Statewide Landcover and Trees Study (SLATS) report, April 1999. pp. 44.

Department of Natural Resources (1999B) Land Cover Change in Queensland 1991-1995, A Statewide Landcover and Trees Study (SLATS) report, July 1999. pp. 52.

Department of Natural Resources (1999C) Land Cover Change in Queensland 1995-1997, A Statewide Landcover and Trees Study (SLATS) report, August 1999. pp. 54.

Department of Natural Resources (2000) Land Cover Change in Queensland 1997-1999, A Statewide Landcover and Trees Study (SLATS) report, September 2000. pp. 56.

Natural Resources and Mines (2002A) Land Cover Change in the Queensland Murray Darling Basin 1999 to 2001, A Statewide Landcover and Trees Study (SLATS) report, July 2002.

Natural Resources and Mines (2002B) Land Cover Change in the Fitzroy Catchment 1999 to 2001, A Statewide Landcover and Trees Study (SLATS) report, October 2002.

Goulevitch, B.M., Danaher, T.J., Stewart, A.J., Harris, D.P., and Lawrence, L.J. (2002) Mapping Woody Vegetation Cover over the State of Queensland using Landsat TM and ETM+ Imagery. In: *Proceedings of the 11th Australasian Remote Sensing and Photogrammetry Conference*, Brisbane, September 2 - 6, 2002.

Henry, B.K., Danaher, T.J., McKeon, G.M. and Burrows, W.H. (2002) A Review of the Potential Role of Greenhouse Gas Abatement in Native Vegetation Management in Queensland's Rangelands. *Rangeland Journal*, 24(1) 2002, 112-32.

Intergovernmental Panel on Climate Change (1996) Greenhouse Gas Inventory Reference Manual, *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories*, Volume 3.

Kuhnell, C., Goulevitch, B., Danaher, T. and Harris, D. (1998) Mapping Woody Vegetation Cover over the State of Queensland using Landsat TM Imagery. In: *Proceedings of the 9th Australasian Remote Sensing and Photogrammetry Conference*, Sydney, Australia, July 1998.

National Forest Inventory (1998) *Australia's State of the Forests Report 1998*. Bureau of Rural Sciences, Canberra.

NGGIC (1998) Australian Methodology for the Estimation of Greenhouse Gas Emissions and Sinks, Workbook for Carbon Dioxide from the Biosphere. National Greenhouse Gas Inventory Committee, Workbook 4.2 Reprinted Revision 2 with Supplements 1998, Canberra.

Tickle, P., Witte, C., Danaher, T. and Jones, K. (1998) The Application of Large-Scale Video and Laser Altimetry to Forest Inventory, In: *Proceedings of the 9th Australasian Remote Sensing and Photogrammetry Conference*, Sydney, Australia, July 1998.

Wedderburn-Bisshop, G.R., Walls, J., Senarath, U., and Stewart, A.J. (2002) Methodology for mapping change in woody landcover over Queensland from 1999 to 2001 using Landsat ETM+. In: *Proceedings of the 11th Australasian Remote Sensing and Photogrammetry Conference*, Brisbane, September 2 - 6, 2002.

Weston, E.J., Harbison, J., Leslie, J.K., Rosenthal, K.M. and Mayer, R.J. (1981) Assessment of the Agricultural and Pastoral Potential of Queensland, Technical Report No. 27, Queensland Department of Primary Industries Agriculture Branch, Brisbane.