

Clearing by Subcatchment

Table 3. Clearing Rates in the SEQ Natural Resource Management Region by subcatchment #.

Period	Clearing rate (,000ha/yr)						
	Albert River	Bribie Island	Brisbane River	Caboolture River	Coomera River	Logan River	Maroochy River
1988–1991	0.067	0.038	0.471	0.785	0.496	0.937	1.005
1991–1995	0.077	0.117	0.385	0.636	0.228	0.598	2.026
1995–1997	0.096	0.185	0.363	0.622	0.311	0.762	1.585
1997–1999	0.086	0.807	0.408	0.303	0.379	0.730	1.093
1999–2000	0.111	0.303	0.577	0.719	0.488	0.777	1.684
2000–2001	0.109	1.063	0.184	0.535	0.279	0.349	1.034
2001–2002	0.321	0.261	0.423	0.232	0.605	0.633	1.801
2002–2003	0.330	0.041	0.945	0.657	1.080	1.432	1.732

Table 3. continued

Period	Clearing rate (,000ha/yr)					
	Moreton Island	Noosa River	North Pine River	South Pine River	Stradbroke Islands	Total
1988–1991	0.000	0.162	0.319	0.199	0.126	4.605
1991–1995	0.000	0.208	0.307	0.132	0.146	4.859
1995–1997	0.000	0.532	0.221	0.096	0.154	4.928
1997–1999	0.005	0.173	0.115	0.135	0.119	4.353
1999–2000	0.000	0.125	0.222	0.121	0.269	5.397
2000–2001	0.000	0.231	0.224	0.076	0.042	4.124
2001–2002	0.002	0.258	0.139	0.051	0.143	4.869
2002–2003	0.002	0.389	0.317	0.152	0.155	7.231




Figure 4. Clearing Rates in the SEQ Natural Resource Management Region by subcatchment #.

Please refer to Figure 21 of the main report (NRM, 2005) for the location of each subcatchment.

Remaining Woody Vegetation

Table 4. Remaining Woody Vegetation in the SEQ Natural Resource Management Region by subcatchment.

	Albert River	Bribie Island	Brisbane River	Caboolture River	Coomera River	Logan River	Maroochy River
Catchment area	78	15	97	47	130	337	139
Area woody 2003	44	11	47	19	75	158	82
% of catchment woody in 2003	57	74	49	41	58	47	59